Journal of Scientific Research and Reports



Volume 30, Issue 8, Page 251-262, 2024; Article no.JSRR.119357 ISSN: 2320-0227

# Effect of Sowing Dates and Nitrogen Levels on Growth, Yield, Quality and Economics of Fodder Maize (Zea mays L.) SFM 1

## Zahida Rashid <sup>a\*</sup>, Saima Nissar <sup>b</sup>, Tanveer Ahmad Ahngar <sup>b</sup>, Raies A Bhat <sup>c</sup>, Sabina Nasseer <sup>a</sup>, Shabeena Majid <sup>a</sup>, Aamir H mir <sup>a</sup>, Sabiya Bashir <sup>d</sup>, Efath Shehnaz <sup>a</sup>, Rakhshanda Anayat <sup>e</sup> and Z. A Dar <sup>a</sup>

<sup>a</sup> Dryland Agriculture Research Station, Rangreth, Srinagar, SKUAST-K, Jammu and Kashmir, India.
<sup>b</sup> Division of Agronomy, Faculty of Agriculture, Wadura, SKUAST-K, Jammu and Kashmir, India.
<sup>c</sup> KVK, Kupwara, Jammu and Kashmir, India.

<sup>d</sup> Division of Plant pathology, Faculty of Horticulture, SKUAST-K, Shalimar, Jammu and Kashmir, India.

<sup>e</sup> Division of Vegetable Science, Faculty of Agriculture, Wadura, SKUAST-K, Jammu and Kashmir, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: https://doi.org/10.9734/jsrr/2024/v30i82245

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/119357

> Received: 05/05/2024 Accepted: 08/07/2024 Published: 25/07/2024

**Original Research Article** 

\*Corresponding author: E-mail: zahida1926@gmail.com;

*Cite as:* Rashid, Zahida, Saima Nissar, Tanveer Ahmad Ahngar, Raies A Bhat, Sabina Nasseer, Shabeena Majid, Aamir H mir, Sabiya Bashir, Efath Shehnaz, Rakhshanda Anayat, and Z. A Dar. 2024. "Effect of Sowing Dates and Nitrogen Levels on Growth, Yield, Quality and Economics of Fodder Maize (Zea Mays L.) SFM 1". Journal of Scientific Research and Reports 30 (8):251-62. https://doi.org/10.9734/jsrr/2024/v30i82245.

## ABSTRACT

The study aimed to assess the effect of sowing dates and nitrogen levels on yield, yield attributes quality, and economic return of Shalimar Fodder Maize-1 (Zea mays L.). An experiment was carried out at Crop Research Farm of the Division of Agronomy, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir, Wadura, during Kharif 2022. The experiment consisted of four sowing dates, S1 (15th April), S2 (15th May), S3 (15th June), and S4 (15th July) and four nitrogen levels, N<sub>0</sub> (0 kg ha<sup>-1</sup>), N<sub>1</sub> (60 kg ha<sup>-1</sup>), N<sub>2</sub> (120 kg ha<sup>-1</sup>) and N<sub>3</sub> (180 kg ha<sup>-1</sup>) as sub-plot treatments, laid out in a split-plot design with three replications. The results showed that the early date of sowing S1 (15th April) had significantly higher values in yield, yield attributes, and quality parameters viz., number of leaves per plant, dry matter accumulation, green fodder yield, dry fodder yield, crude protein content (%), ADF (%), NDF (%), and ash content (%) than other sowing dates, while S<sub>4</sub> (15th July) had significantly lower growth, yield, and guality parameters. However, S<sub>1</sub> (15<sup>th</sup> April) recorded a significantly lower leaf stem ratio, which was seen to be decreasing with an increase in sowing dates. Among different nitrogen levels, the yield and quality parameters viz. leaf stem ratio, number of leaves per plant, dry matter accumulation, green fodder yield, dry fodder yield, crude protein, and ash content were significantly influenced by increasing levels of nitrogen, being maximum with N<sub>3</sub> (180 kg ha<sup>-1</sup>), which was at par with N<sub>2</sub> (120 kg ha<sup>-1</sup>), whereas N<sub>0</sub> (control) recorded significantly lower growth, yield, and quality parameters. Furthermore, NDF (%) and ADF (%) decreased as nitrogen levels increased, and were lowest at  $N_3$  (180 kg ha<sup>-1</sup>). In terms of economic suitability, the treatment combination of  $S_1N_3$  (15<sup>th</sup> April + 180 kg ha<sup>-1</sup>) had the highest net returns (Rs. 143841.7 ha<sup>-1</sup>) and benefit: cost ratio (2.87), followed by  $S_1N_2$  (15<sup>th</sup> April + 120 kg ha<sup>-1</sup>) with net return (Rs. 120721.3 ha<sup>-1</sup>) and benefit: cost ratio (2.45).

Keywords: Fodder maize; sowing dates; nitrogen levels; yield, quality; economic return; grain basket, maize.

## **1. INTRODUCTION**

"Maize crop has an important place in the food grain basket of our country and is the third most important food crop because of its importance in food, feed, fodder, speciality corn, starch etc. It has yield potential far higher than any other cereal hence referred to as the miracle crop or the "Queen of Cereals". In the world, nearly 1147.7 million tonnes of maize are being produced together by over 170 countries from an area of 193.7 million hectares with average productivity of 5.75 t ha-1" [1]. "It has attained a position of industrial crop globally as 83% of its production in the world is used in feed. starch and biofuel industries. The global consumption pattern of maize is feed 61%, food 17% and industry 22%. It is a prime driver of the global agricultural economy" [1]. "India ranks fourth in area and seventh in production among maizegrowing countries, encompassing around 4% of global maize area and 2% of total production. During 2018-2019 in India, the maize area has reached to 9.2 million hectares" [2]. "During 1950-51, India had only 1.73 million MT of maize, which has increased to 27.8 million MT, representing an almost 16-fold increase in production. The average productivity during the period has increased by 5.42 times from 547 kg

ha<sup>-1</sup> to 2965 kg ha<sup>-1</sup>, while the area increased nearly by three times. Maize is the second most important cereal crop after rice in the Union Territory of Jammu and Kashmir, grown on an area of 0.31 million hectares with a production of 0.51 million tonnes and an average productivity of 1650 kg ha-1" [3]. "In the union territory of Jammu and Kashmir, about 90.6% of the total area under maize is planted under rainfed conditions" [4]. "India ranks fourth in area and seventh in production among maize-growing countries, encompassing around 4% of global maize area and 2% of total production. During 2018-2019 in India, the maize area has reached to 9.2 million hectares" [2]. "During 1950-51, India produced only 1.73 million MT of maize, which has increased to 27.8 million MT, representing an almost 16-fold increase in The production. average productivity during the period has increased by 5.42 times from 547 kg ha<sup>-1</sup> to 2965 kg ha<sup>-1</sup>, while the area increased nearly by three times. Maize is the second most important cereal crop in the union territory of Jammu and Kashmir, after rice, and is grown on an area of 0.31 million hectares, with a production of 0.51 million tonnes and an average productivity of 1650 kg ha-1" [3]. "In the union territory of Jammu & Kashmir, about 90.6% of the total area

under maize is grown under rainfed circumstances" [4].

"Maize (Zea mays L.) is considered an important double-purpose crop (used in the human diet as grain as well as animal feed) grown widely in the kharif season. Maize is generally sown under rainfed conditions and in marginal areas, particularly in hilly terrains of the Kashmir valley invariably as an intercrop with pulses. Maize requires considerable moisture and temperature from germination to flowering. The ideal temperature for germination is 21°C, whereas for growth it is 32°C. Maize is quite sensitive to stagnant water, especially in the early phases of growth" (Chaudhary et al., 2012).

"Maize forage supplies a substantial amount of energy-rich forage for dairy animal diets, free from anti-nutritional factors unlike sorahum so can be fed to cattle at any growth stage" [5]. "Forage maize has recently become a crucial component of ruminant rations, as it enriches diets and improves dairy COW animal performance" [6,7]. "The maize crop has many desirable fodder qualities, such as high productivity, high production potential, broader adaptability, rapid growing nature, succulence, palatability, toxin-free, rich in protein and minerals, and high digestibility, which makes it preferred over other cultivated non-fodder crops, resulting in increased body weight and milk quality in cattle" [8]. "Hence it is widely known as a ready-made fodder crop. Maize is highly nutritive, excellent and sustainable fodder for livestock" [9]. "The green fodder of maize possesses lactogenic properties and therefore, is suited for milch cattle" [10]. Maize possesses high soluble sugar concentrations in the green stage, making it suitable for silage preservation, and it is referred to as the king of crops for silage-making.

"Sowing time of maize varies from place to place, especially at different altitudes in Kashmir. Sowing time of maize in lower belts of Kashmir ranges from last week of April to mid of May. There is the modifying effect on the growth and development of maize plants with the environmental changes (sunshine, temperature) associated with different sowing dates. Sowing dates had a significant effect on maize growth and yield. Greater the deviation from the optimal maize sowing date, higher the reduction in yield" [11,12]. "One of the challenges for maize grower's present is to find the narrow interval

between sowing earlier and sowing later" [13]. "Lower yield is obtained when seeds are sown too early or late, probably due to the presence of unfavourable climatic conditions before or after the growing season. Maize sown earlier performs better and produces better yields due to the fact that the vegetative phase of the life cycle occurs during the colder part of the season with less moisture stress" [14]. "Late sowing reduces maize production by shortening the effective growing season, reducing crop photosynthesis, and exposing the crop to high-risk lethal cold temperatures before grain maturation, insect and disease stress, and moisture and heat stress during pollination" [15]. "Sowing time is such a changeable element of agro-technique which does not require extra cost however, we can manage it to increase the efficiency of crop. In addition to recommended agro-management measures, choosing an appropriate sowing date with a good variety guarantees higher maize production" [16]. "For achieving higher efficiency from available nutrients, water and solar energy. the selection of optimum sowing date for already existing varieties is unavoidable" [17].

"Fertilizer application is one of the principal factors that directly influence fodder yield and quality. An adequate supply of nutrients at each growth stage is highly essential for improving yield and quality of fodder maize. Nitrogen is the most important essential element for plants and is required in a comparatively large amount. It is the basic constituent of chlorophyll, protoplasm, amino acids and nucleic acids. It enhances the growth and development of all living tissues and helps in improving the quality of fodder and the protein content of grains. It is the most important element vield-limiting factor in agricultural systems" [18]. It is mainly supplied through mineral fertilizers, farm manure. symbiotic nitrogen fixation, and atmospheric wet and dry deposition. It is very essential for plant growth and makes up 1 to 4% of plant dry matter. It also mediates the utilization of phosphorus, potassium, and other elements in plants.

Therefore, a field investigation was conducted to evaluate the effect of dates of sowing and nitrogen levels on yield attributes, yield, quality and economics of fodder maize (*Zea mays* L.) SFM-1 under conditions of Kashmir valley.

## 2. MATERIALS AND METHODS

The experiment was conducted at the Crop Research Farm of Division of Agronomy, Faculty

of Agriculture Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir, Wadura Sopore during Kharif, 2018. The site is located between 34º 21' N latitude and 74º 23'E longitude and has an altitude of 1590 m above mean sea level. The experimental site was welldrained with uniform topography. The experimental site falls in a mid to high altitude characterized by a temperate climate with hot summers and severe cold winters. The average annual precipitation is 812 mm most of which is received from December to April in the form of snow and rains. The soil of the experimental field was silt clay loam in texture with slightly acidic to neutral pH, medium in organic carbon, normal electrical conductivity and the available nitrogen, phosphorus and potassium were also found medium.

The experiment comprised of factors with four sowing dates *viz.*,  $S_1$  (15<sup>th</sup> April),  $S_2$  (15<sup>th</sup> May),  $S_3$  (15<sup>th</sup> June) and  $S_4$  (15<sup>th</sup> July) as main plot treatments and four nitrogen levels *viz.*,  $N_0$  (0 kg ha<sup>-1</sup>),  $N_1$  (60 kg ha<sup>-1</sup>),  $N_2$  (120 kg ha<sup>-1</sup>) and  $N_3$  (180 kg ha<sup>-1</sup>) as sub-plot treatments laid out in split plot design with three replications. The total number of treatment combinations was 16 and the total number of treatment plots was 48. The gross plot size of each subplot was 15.8 m<sup>2</sup>.

Leaf stem ratio: Five plants from each plot were taken from the base and the leaves were separated from the stems of each plant for the leaf stem ratio. The leaves and stems from each plot were sun-dried and weighed individually. The leaf to stem ratio was then computed as

Leaf stem ratio  $= \frac{Dry \ weight \ of \ leaves}{Dry \ weight \ of \ stem}$ 

**Number of leaves plant**<sup>-1</sup>: Five plants were selected randomly to count the number of leaves. The number of green leaves considered as leaves plant<sup>-1</sup> was recorded at 30 DAS, 60 DAS, and harvest.

**Dry matter accumulation:** Plant samples were randomly selected from the penultimate rows of each plot at 30 DAS, 60 DAS, and harvest. The samples were cut from the base with a knife when they were small, and then with a sickle when they grew bigger. These samples were dried in the oven for 4-6 days to obtain a constant weight at temperatures ranging from 60-

 $65^{\circ}$ C. The weight was recorded in grams and then converted to q ha<sup>-1</sup>.

**Green fodder yield:** The green fodder yield of each net plot (leaving the border and penultimate rows) was harvested, and the weight from each plot was recorded individually as kg plot<sup>-1</sup>, which was further converted into q ha<sup>-1</sup>.

**Dry fodder yield:** The fresh fodder yield of each net plot was allowed to sundry for about a week and weight was recorded in kg and converted to q ha<sup>-1</sup> to determine the dry weight ha-1.

**Crude protein:** The nitrogen (N) concentration was assessed by using the modified micro-Kjeldahl technique and expressed in percentage. Protein content was estimated by multiplying the N content by a factor 6.25.

Ash content: The ash content of ground samples was evaluated using the method described by AOAC, [19] The following formula was used to calculate the total ash, which was then expressed in percentages.

Total ash (%) = 
$$\frac{Weight of ash}{Weight of sample} X 100$$

**ADF and NDF (%):** The concentrations of neutral detergent fiber (NDF) and acid detergent fiber (ADF) were measured using proximate analysis Van Soest fiber analysis, Goering and Soest, [20] When a sample is boiled (refluxed) in a neutral detergent solution for an hour, all the cell contents dissolve completely. The residue left after boiling is known as cell wall content. To determine the percentage of cell wall content in a sample, the cell wall content is dried overnight at 100-105°C and then weighed. Cell content is determined by the difference, *i.e.*, % cell content = 100 - % cell wall content.

Calculation:

Weight of sample = W<sub>1</sub> g Weight of empty crucible = W<sub>2</sub> g Weight of crucible + NDF = W<sub>3</sub> g Weight of NDF = (W<sub>3</sub>-W<sub>2</sub>) g (% of NDF) =  $\frac{W_3 - W_2 X 100}{W_1}$  When the sample is boiled in an acid detergent solution for an hour, all the cell content and hemicellulose dissolve completely. The residue left after boiling is known as acid detergent fiber. The acid detergent fiber (ADF) is then dried and washed to find out the ADF % in the sample. Calculation:

Weight of sample =  $W_1 g$ 

Weight of empty crucible =  $W_2 g$ 

Weight of crucible + ADF =  $W_3 g$ 

Weight of ADF =  $(W_3-W_2)$  g

$$(\% of ADF) = \frac{W_3 - W_2 X 100}{W_1}$$

**Relative economics:** The economics of various treatments were calculated using the cost of cultivation, gross return, net return, and benefit-cost ratio. The cost of cultivation was calculated for each treatment based on the price of input. The gross return was calculated based on the existing price of the output. The benefit-cost ratio was then determined to estimate the profitability of the per-rupee investment.

Net returns = Gross returns - Total cost of cultivation

Benefit: cost ratio = Net returns ÷ Total cost of cultivation

## 3. RESULTS AND DISCUSSION

## 3.1 Effect of Sowing Dates

#### 3.1.1 Growth and Yield

Data showed that sowing dates have a significant effect on leaf stem ratios taken at 30day intervals. A perusal of the data in Table 1 showed that the leaf stem ratio was higher on the 15<sup>th</sup> July (S<sub>4</sub>) sowing than the rest of the sowings up to 30 DAS. However, at 30 DAS, the sowing date of 15<sup>th</sup> April (S<sub>1</sub>) had a higher leaf stem ratio (0.51 cm) than the rest of the sowing dates. The highest leaf: stem ratio was obtained on the 15th of April (S<sub>1</sub>) compared to other sowing dates, which was mostly due to more stem weight and less leaf weight in comparison to the other sowing dates. Saimaheshwari et al. (2020) found that "delay in sowing significantly decreased leaf stem ratio. However, up to 30 DAS, 15th July (S<sub>4</sub>) exhibited highest the leaf stem ratio

compared to the other sowing dates and is contributed by more stem weight and more leaf weight initially".

The data in Table 1 showed that the number of leaves plant<sup>-1</sup> was significantly affected by the sowing dates recorded at different stages. Sowing on 15<sup>th</sup> April (S<sub>1</sub>) resulted in the highest number of leaves per plant (14.02) while sowing on July 15<sup>th</sup> (S<sub>4</sub>) obtained the lowest number of leaves per plant. It might be due to the presence of ideal growing conditions at the time of sowing. "The number of leaves per plant decreased with delayed sowing due to a reduction in photoperiod" [21]. The results are comparable to the findings of Beiragi et al. [22], who discovered that "delayed planting decreased the total number of leaves per plant". Similarly, Martin and Williams (2008) discovered that "delayed sowing results in a decrease in leaf number per plant and slower rates of leaf appearance".

The data on dry matter accumulation presented in Table 1 demonstrated that the sowing date significant effect on had а drv matter accumulation. "Sowing date of 15th April recorded the highest matter (S<sub>1</sub>) dry accumulation than the rest of the sowing dates. The reason might be attributed to the exploitation of favourable climatic conditions at important stages of growth by the 15<sup>th</sup> April (S<sub>1</sub>) sown crop, and higher leaf number, which might have provided more photosynthetic area (LAI) and contributed more dry matter. The lowest dry matter accumulation was observed on July 15th (S<sub>4</sub>) sowing may be due to unfavourable growth conditions" [21]. Girijesh et al. [23] revealed that "earlier and timely sowing significantly increases dry matter production in maize crops due to the adequate growing period available during the growth phase, compared to delayed sowing". Cirilo and Andrade [24] and Maddonni et al. [25] found that delayed maize sowing reduces dry matter yield. However, the 15th of July (S4) sowing exhibits a higher dry matter content up to 30 DAS than other sowing dates due to temperatures, considerably greater which accelerated growth initially.

The data regarding green fodder yield is presented in Table 2 which indicates that green fodder yield was significantly influenced by sowing dates. Sowing on April  $15^{th}$  (S<sub>1</sub>) resulted in the highest green fodder yield (520.33 q ha<sup>-1</sup>), whereas sowing on July  $15^{th}$  (S<sub>4</sub>) resulted in the lowest green fodder yield (382 q ha<sup>-1</sup>).

Treatments	Leaf Stem Ratio			Number of Leaves Plant <sup>-1</sup>			Dry Matter Accumulation (q ha <sup>-1</sup> )		
	30 DAS	60 DAS	Harvest	30 DAS	60 DAS	Harvest	30 DAS	60 DAS	Harvest
Main Plot Dates of so	owing								
S <sub>1</sub> (15 <sup>th</sup> April)	0.99	0.89	0.51	5.64	12.15	14.02	23.20	106.70	162.07
S <sub>2</sub> (15 <sup>th</sup> May)	1.12	0.80	0.44	6.14	11.42	12.85	27.18	99.85	147.73
S <sub>3</sub> (15 <sup>th</sup> June)	1.21	0.71	0.37	6.65	10.69	11.84	30.81	93.89	136.53
S <sub>4</sub> (15 <sup>th</sup> July )	1.30	0.61	0.30	7.15	9.97	10.83	33.23	88.49	127.75
SEm±	0.02	0.02	0.01	0.14	0.19	0.28	1.02	1.58	2.19
CD (p≤0.05)	0.08	0.08	0.06	0.49	0.70	1.00	3.62	5.59	7.75
Sub Plot Nitrogen lev	/els								
N <sub>0</sub> (0 kg ha <sup>-1</sup> )	1.07	0.61	0.32	4.13	7.23	8.45	16.15	82.00	124.38
N <sub>1</sub> (60 kg ha <sup>-1</sup> )	1.13	0.73	0.38	6.49	11.72	13.04	27.65	96.47	142.13
N <sub>2</sub> (120 kg ha <sup>-1</sup> )	1.19	0.81	0.44	7.19	12.38	13.73	33.75	102.88	150.36
N₃ (180 kg ha⁻1)	1.23	0.86	0.48	7.76	12.90	14.33	36.87	107.57	157.21
SEm±	0.01	0.02	0.01	0.19	0.21	0.23	1.14	1.61	2.33
CD (p≤0.05)	0.05	0.07	0.05	0.58	0.65	0.68	3.35	4.74	6.86

## Table 1. Effect of sowing dates and nitrogen levels growth of Shalimar Fodder Maize-1

Table 2. Effect of sowing dates and nitrogen levels on yield of Shalimar Fodder Maize-1

Treatments	Green Fodder Yield (q ha <sup>-1</sup> )	Dry Fodder Yield (q ha <sup>-1</sup> )
Main Plot Dates of Sowing		
S <sub>1</sub> (15 <sup>th</sup> April)	520.33	208.13
S <sub>2</sub> (15 <sup>th</sup> May)	489.22	195.68
S₃ (15 <sup>th</sup> June )	441.44	176.57
S <sub>4</sub> (15 <sup>th</sup> July )	382.00	152.80
SEm±	9.03	3.61
CD (p≤0.05)	31.05	12.34
Sub Plot Nitrogen Levels		
N <sub>0</sub> (0 kg ha <sup>-1</sup> )	350.25	140.104
N1 (60 kg ha <sup>-1</sup> )	430.22	172.08
N <sub>2</sub> (120 kg ha <sup>-1</sup> )	506.44	202.57
N₃ (180 kg ha <sup>-1</sup> )	546.07	218.42
SEm±	13.53	5.41
CD (p≤0.05)	39.75	15.90

Crops sown earlier got more time to utilize available resources and favourable temperature at later growth stages of the crop, whereas late sown crops got less time to utilize growth factors, which is responsible for lower leaf area index and stunted plant growth, which decreases dry matter production and insufficient assimilate partitioning towards sink for superior vegetative growth, resulting in a reduction of yield and yield contributing elements in comparison to timely sown crop. Similar findings were reported by Varma et al. [26] and Shargi et al. [27].

The data on dry fodder yield in Table 2 indicated that sowing date had a significant effect on dry fodder yield. Among sowing dates, 15<sup>th</sup> April (S<sub>1</sub>) recorded the highest dry fodder yield (208.12 q ha<sup>-1</sup>), while 15th July (S<sub>4</sub>) recorded the lowest dry fodder yield (152.80 q ha-1). A decrease in dry fodder yield was recorded with successive delays in sowing time. Lower fodder yield in late-sown forage maize could be due to less forage production as a result of lower temperatures at the early growth stages in late-sown crops compared to early-sowed crops. Dar et al. [28] similarly found that early sowing resulted in higher forage yield than late sowing, which they attributed the available congenial to temperature for better growth and development of forage maize.

#### 3.1.2 Quality parameters

The data regarding crude protein content is given in Table 3 which indicates sowing dates had a statistically non-significant effect on crude protein. The highest protein content (9.23%) was obtained on April 15th (S1), while the lowest protein content (8.90%) was recorded on July 15<sup>th</sup> (S<sub>4</sub>) sowing date. Sowing fodder maize on April 15<sup>th</sup> (S<sub>1</sub>) registered the highest protein content, whereas sowing on July 15th (S<sub>4</sub>) showed the lowest crude protein content. Since protein content is inversely proportionate to green fodder yield and the genetic character of the plant remained non-significant among all the dates of sowing. Darby and Lauer [29] observed that delayed sowing results in low forage quality. Mokhtarpur et al. [30] discovered that delayed sowing had an inverse effect on protein content, and the highest protein content was observed in early sowing.

The data presented in Table 3 indicated that neutral detergent fiber and acid detergent fiber differed significantly with respect to sowing

dates. Sowing on 15th April (S1) reported significantly higher neutral detergent fiber (69.23%) and acid detergent fiber (46.98) respectively. While the lowest neutral detergent fiber (62.61%) and acid detergent fiber (38.98%) were determined on 15<sup>th</sup> July (S<sub>4</sub>) sowing. "Early sowing increases (NDF%) and (ADF%) because the crop sown earlier attains full maturity because of the favourable temperature during the growing period of 1<sup>st</sup> sowing, the crop attained the maximum plant height, plant dry weight which favours the maximum ADF and NDF resulting in more accumulation of lignin, cellulose and hemicellulose in forage than late sowing in which the crop does not attain full maturity thus resulting in lower neutral detergent fiber (NDF%) and acid detergent fiber (ADF%) in forage" [21] Salama [31] discovered that "neutral detergent fiber and acid detergent fiber are dependent on the age of the plant at harvest and increase with maturity, so the age of plants at harvest varies with different sowing dates, which consequently increases or decreases the neutral detergent fiber and acid detergent fiber in forage". Similar conclusions were drawn by Gaile [32] and Ball et al. [33].

The data on ash content in Table 3 demonstrated that sowing dates had a significant effect on ash content. The highest ash level (8.73%) was observed on the  $15^{th}$  of April (S<sub>1</sub>). The sowing date of the  $15^{th}$  of July (S<sub>4</sub>) resulted in the lowest ash content (8.31%). The highest ash content on April  $15^{th}$  (S1) may be attributed to increased dry matter content, which improved the nitrogen uptake by the plants. Koca et al. [34] revealed that early sowing results in higher ash content than late sowing.

A perusal of the data given in Table 1 showed that nitrogen levels had a significant effect on leaf: stem ratio. Among the various nitrogen levels. N<sub>3</sub> (180 kg Ν ha<sup>-1</sup>) recorded significantly higher leaf stem ratios at 30 DAS (1.23 cm), 60 DAS (0.86 cm), and harvest (0.48 cm) compared to  $N_1$  (60 kg N ha<sup>-1</sup>) and  $N_0$ (Control). However, it was at par with N<sub>2</sub> (120 kg N ha-1). The lowest leaf: stem ratio was observed in N<sub>0</sub> (Control). It was mainly due to the rapid expansion of dark green foliage, which could intercept and utilize the incident solar energy in the production of photosynthates, resulting in higher meristematic activity and leaf: stem ratio of fodder maize. Manjanagouda et al. [35] and Somashekhar [36] reported that an increase in nitrogen levels increased the leaf:

Treatments	Crude Protein (%)	Ash Content (%)	Neutral Detergent fiber (NDF %)	Acid Detergent Fiber (ADF %)		
Main Plot Dates of	f Sowing					
S <sub>1</sub> (15 <sup>th</sup> April)	9.23	8.73	69.23	46.98		
S <sub>2</sub> (15 <sup>th</sup> May)	9.04	8.60	67.67	44.77		
S <sub>3</sub> (15 <sup>th</sup> June )	8.97	8.43	65.15	41.57		
S₄ (15 <sup>th</sup> July )	8.90	8.31	62.61	39.98		
SEm±	0.17	0.16	0.23	0.26		
CD (p≤0.05)	NS	NS	0.82	0.92		
Sub Plot Nitrogen Levels						
N <sub>0</sub> (0 kg ha <sup>-1</sup> )	8.39	7.69	70.60	48.01		
N1 (60 kg ha <sup>-1</sup> )	8.83	8.42	66.50	44.77		
N <sub>2</sub> (120 kg ha <sup>-1</sup> )	9.26	8.80	64.45	41.57		
N₃ (180 kg ha⁻¹)	9.67	9.16	63.12	40.28		
SEm±	0.14	0.12	0.67	0.63		
CD (p≤0.05)	0.42	0.37	1.99	1.86		

#### Table 3. Effect of sowing dates and nitrogen levels on Quality parameters of Shalimar Fodder Maize -1

## Table 4. Relative economics of shalimar fodder maize-1 under varied sowing dates and nitrogen levels

Treatments	CCost of	Gross	Net Returns	B:C
	Cultivation (₹ ha <sup>-1</sup> )	Returns (₹ ha⁻¹)	(₹ ha⁻¹)	Ratio
S1N0	47690	116100	68410	1.43
S1N1	48299	144400	96101	1.99
$S_1N_2$	49212	169933.3	120721.3	2.45
<b>S</b> 1 <b>N</b> 3	50125	193966.7	143841.7	2.87
$S_2N_0$	47690	112100	64410	1.35
$S_2N_1$	48299	137733.3	89434.33	1.85
$S_2N_2$	49212	159933.3	110721.3	2.25
S <sub>2</sub> N <sub>3</sub>	50125	177300	127175	2.54
$S_3N_0$	47690	102100	54410	1.14
S <sub>3</sub> N <sub>1</sub>	48299	124400	76101	1.58
S <sub>3</sub> N <sub>2</sub>	49212	145933.3	96721.33	1.97
S <sub>3</sub> N <sub>3</sub>	50125	157300	107175	2.14
$S_4N_0$	47690	90010	42320	0.89
S <sub>4</sub> N <sub>1</sub>	48299	109733.3	61434.33	1.27
$S_4N_2$	49212	131933.3	82721.33	1.68
S <sub>4</sub> N <sub>3</sub>	50125	126723.3	76598.33	1.53

stem ratio. The data regarding the number of leaves plant<sup>-1</sup> is given in Table 1 which indicates that the number of leaves plant<sup>-1</sup> in fodder maize was significantly influenced by nitrogen levels. Among N levels, N<sub>3</sub> (180 kg N ha<sup>-1</sup>) resulted in a higher number of leaves, however, it was at par with  $N_2$  (120 kg ha<sup>-1</sup>). The  $N_0$  (Control) recorded the lowest number of leaves. Mahdi et al. [37] also observed "a significant increase in the leaf number plant-1 with an increase in nitrogen levels. An increase in the number of green leaves might be attributed to the availability of sufficient nitrogen at all stages of growth". "Plant height increased the number of nodes plant<sup>-1</sup>, resulting in more number of leaves plant<sup>-1</sup>. Again, nitrogen helped in the rapid growth

and development by helping in photosynthesis and various biochemical processes which respond to growth" [38].

The data in Table 1 showed that sowing dates had a significant impact on dry matter accumulation. Among nitrogen levels, N<sub>3</sub> (180 kg N ha<sup>-1</sup>) recorded the highest dry matter accumulation at 30 DAS (36.87 q ha<sup>-1</sup>), 60 DAS (107.57 q ha<sup>-1</sup>) and at harvest (157.21 q ha<sup>-1</sup>), which was at par with N<sub>2</sub> (120 kg N ha<sup>-1</sup>). However, N<sub>0</sub> (Control) recorded the lowest dry matter accumulation. The increase in dry matter production at higher nitrogen levels might be due to enhanced metabolic activity as well as increased cell growth and elongation. An

increased dose of Nitrogen enhanced the photosynthetic surface area (LAI), resulting in higher dry matter accumulation. The findings are in accordance with the findings revealed by Puri and Tiwana (2008), Ali et al. [39] and Mahdi et al. [37].

Data presented in Table 2 indicated that green fodder yield was significantly influenced both by nitrogen levels and among nitrogen levels N<sub>3</sub> (180 kg N ha<sup>-1</sup>) recorded maximum green fodder yield (546.07 q ha  $^{-1}$ ) whereas N<sub>0</sub> (Control) recorded significantly lowest green fodder yield (350.25 g ha<sup>-1</sup>). However, N<sub>3</sub> (180 kg N ha<sup>-1</sup>) was found to be statistically at par with N2 (120 kg N ha-1). The highest green fodder yield with the N<sub>3</sub> level is due to efficient growth and yield attributes such as plant height, leaf area, number of leaves per plant and dry matter production. Nitrogen is an essential component of plant tissue and is associated with cell division and elongation. Therefore its positive impact on growth parameters resulted in better vield. These results are in accordance with the findings of Panwar et al. [40], who observed that increasing nitrogen levels significantly increases green fodder yield.

The data regarding dry fodder yield is presented in Table 2 which indicates that dry yield was significantly influenced by nitrogen levels. Among nitrogen levels, N<sub>3</sub> (180 kg N ha<sup>-1</sup>) recorded the highest dry fodder yield (218.42 g ha<sup>-1</sup>), whereas N<sub>0</sub> (Control) recorded the lowest. However, N<sub>3</sub> (180 kg N ha<sup>-1</sup>) was statistically at par with N<sub>2</sub> (120 kg N ha-1). Since nitrogen is an important component of plant tissue and plays an essential role in cell division and elongation, its beneficial effect on growth characteristics such as plant height, leaf area index, and stem diameter might have contributed to increased yield. Eltelib et al. [41] also reported similar results. The increase in fodder yield was evidently due to the cumulative effect of increased growth parameters, which ultimately resulted in increased green and dry fodder yields. Budakli et al. [42], Aslam et al. [43], Patel et al. [44], and Meena et al. [45] also reported similar findings.

## 3.1.3 Quality parameters

The data given in Table 3 showed that nitrogen levels had a significant effect on the crude protein content of fodder maize. Among Nitrogen levels,  $N_3$  (180 kg N ha<sup>-1</sup>) recorded the highest protein content, whereas  $N_0$  (Control) recorded the lowest. However,  $N_3$  (180 kg N ha<sup>-1</sup>) was statistically at par with  $N_2$  (120 kg N ha<sup>-1</sup>).

Ullah et al. (2018), Mahdi et al. [37] and Eltelib et al. [41] also confirmed that increasing nitrogen levels increases crude protein content. This might be due to the fact that plants have a large concentration of nitrogen at higher N levels, which in turn accelerates the production of nucleotides and coenzymes for protein synthesis Kakol et al. [46].

A perusal of the data given in Table 3 revealed that neutral detergent fiber and acid detergent fiber differed significantly with respect to nitrogen levels. Among nitrogen levels, both neutral detergent fiber and acid detergent fiber were seen be decreasing as nitrogen levels increased, while N0 (Control) recorded the highest neutral detergent fiber (70.60%) and acid detergent fiber (48.01%) than other nitrogen levels. While the lowest neutral detergent fiber (63.12%) and acid detergent fiber (40.28%) were recorded in  $N_3$  (180 kg N ha<sup>-1</sup>), which was at par with  $N_2$  (120 kg N ha<sup>-1</sup>). The results are in accordance with the findings reported by Almodares et al. [47], who observed that fiber content decreased with an increase in nitrogen levels.

The data in Table 3 revealed sowing dates had a significant impact on ash content. Among nitrogen levels, the highest ash content (9.16%) was recorded in  $N_3$  (180 kg N ha<sup>-1</sup>), which was statistically at par with  $N_2$  (120 kg N ha<sup>-1</sup>), and is attributed to the greater dry matter content, which improved mineral matter. However, the lowest ash content (7.69%) was recorded in  $N_0$  (Control). Saruhan and Sireli [48] also found that increasing the nitrogen dose increased the ash content of fodder maize.

## 3.1.4 Relative economics

The data in Table 4 showed that sowing dates and nitrogen levels had a significant effect on gross returns, net returns, and benefit: cost ratio. The relative economics in terms of gross returns, net returns, and benefit: cost ratio with respect to vield was green fodder worked out for various treatment combinations [49,50]. The data revealed that fodder maize sown on the 15<sup>th</sup> April (S<sub>1</sub>) date of sowing and Nitrogen level N<sub>3</sub> (180 kg N ha<sup>-1</sup>) recorded the highest gross returns (₹ 193967), net returns (₹ 143842), and benefit: cost ratio (2.87) followed by 15<sup>th</sup> May sowing with  $N_2$  (120 kg N ha^-1) [51,52]. The lowest gross returns, net returns and benefit: cost ratio were recorded by fodder maize on 15th June (S1) sowing with Control (N<sub>0</sub>). This could be due to higher green fodder yield in these treatment combinations than in others. The findings are in agreement with Patel et al. [44].

## 4. CONCLUSION

In conclusion, the results of the investigation indicated that fodder maize sown on  $15^{th}$  April (S<sub>1</sub>) and nitrogen level N<sub>2</sub> (120 kg ha<sup>-1</sup>) produced higher yield attributes, yield, quality and economics of fodder maize however, it was statistically at par with sowing done on  $15^{th}$  April (S<sub>1</sub>) and nitrogen level N<sub>2</sub> (120 kg ha<sup>-1</sup>). Highest gross returns, net returns and benefit: cost ratio were recorded with the  $15^{th}$  April (S<sub>1</sub>) sowing date and with a combination of nitrogen level N<sub>3</sub> (180 kg N ha<sup>-1</sup>) followed by N<sub>2</sub> (120 kg ha<sup>-1</sup>).

Based on the above studies we concluded that under Kashmir valley conditions, sowing of fodder maize should be carried around  $15^{th}$  April (S<sub>1</sub>) with nitrogen level N<sub>2</sub> (120 kg ha<sup>-1</sup>).

## **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## ACKNOWLEDGEMENTS

The paper is a part of M. Sc. Thesis of the second author submitted to Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir. I wish to express my profound gratitude to Dr. Zahida Rashid for her combined support, encouragement and suggestions throughout the course and conduct of this work.

## **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

## REFRENCES

- 1. FAOSTAT. Food and Agricultural Organization of the United Nations; 2020. Available:https://iimr.icar.gov.in/worldmaze-scenario/
- 2. DACNET. Department of Agriculture and Cooperation, India; 2020.
- 3. Directorate of Economics and Statistics; 2019.

- 4. Directorate of Economics and Statistics; 2017.
- Dahmardeh M, Ghanbari A, Syasar B, Ramroudi M. Effect of intercropping maize with cowpea on green forage yield and quality evaluation. Asian Journal of Plant Science. 2009; 8(3):235-239.
- Anil L, Park J, Phipps RH. The potential of forage – maize intercrops in ruminant nutrition. Animal Feed Science and Technology. 2000;86(3-4):157-184.
- Cusicanqui JA, Hinds FC, Marchello JA. Effects of dietary forage and protein levels on lactation and digestion in dairy cows. Journal of Dairy Science. 1999;82(6):1276-1289.
- Sattar MA, Haque MF, Rahman MM. Intercropping maize with broadcast rice at different row spacing. Bangladesh Journal of Agricultural Research. 1994;19:159– 164.
- Iqbal A, Ayub M, Zaman H, Ahmed R. Impact of nutrient management and legume association on agro - qualitative traits of maize forage. Pakistan Journal of Botany. 2006;38:1079–1084.
- Valk H. Effect of reducing nitrogen fertilizer on grassland on grass intake, digestibility and milk production of dairy cows. Livestock Production Science. 2000;63(1):27-38.
- 11. Sarvari M, Futo Z. Correlation between the sowing time, yield, and seed moisture content at harvest on chernozem soil. Acta Agraria Debreceniensis. 2000;1(10):32-41.
- 12. Berzsenyi Z, Lap DQ. Effect of sowing time and N fertilisation on the yield and yield stability of maize (*Zea mays* L.) hybrids between 1991-2000. Novenytermeles 2001;50(2):309-331.
- Nielsen RL, Thomson PR, Brown GA, Halter AL, Wells J, Wuethricc KL. Delaying planting date effects on flowering and grain maturation of corn. Agronomy Journal. 2002;93:540-547.
- 14. Ali Q, Ali A, Waseem M, Muzaffar A, Ahmad S, Ali S, Awan M, Samiullah T. Correlation analysis for morphophysiological traits of maize (*Zea mays* L.). Life Science Journal. 2014;11:9-13.
- 15. Otegui ME, Melon S. Kernel set and flower synchrony within the ear of maize. I. Sowing date effects. Crop Science. 1997;37:441-447.
- 16. Qureshi AS, Qadir M, Heydari N, Turral H, Javad A. A review of management strategies for salt-prone land and water

resources in Iran. International water management Institute 30P (IWMI working paper 125) Colombo, Srilanka; 2007.

- Ogbomo KEL, Remison SU. Growth and yield of maize as influenced by sowing date and poultry manure application. Notulea. Botanicae. Horti. Agrobotanici Cluj. 2009;37(1):199-203.
- Subedi, Ma BL. Assessment of major yield limiting factors 0f maize production in humid temperate environment. Field Crop Research. 2009; 110(1):21-26.
- AOAC. Official Methods of Analysis. Association of official analytical chemists, 18<sup>th</sup> edn. Arlington, Virginia, USA; 1995.
- 20. Van-Soest PJ, Robertson JB, Lewis BA. Methods for dietary fiber, neutral detergent fiber and non-starch polysaccharides in relation to animal nutrition. Journal of Dairy Sciences. 1970;74 (10):3583-97.
- Nisar S, Rashid Z, Touseef A, Kumar R, Nissa SU, Faheem J, Angrez A, Sabina N, Shabeena M, Tanveer A, Amal S. Productivity of fodder maize (*Zea mays* L.) SFM-1 under varied sowing dates and nitrogen levels. International Journal of Bio-resource and Stress Management. 2024;15:01-12.
- Beiragi MA, Khorasani SK, Shojaei SH, Dadresan M, Mostafavi K, Golbashy M. A study on effects of planting dates on growth and yield of 18 corn hybrids (*Zea* mays L.). American Journal of Experimental Agriculture. 2011;1(3):110-120.
- 23. Girijesh GK, Swamy ASK, Sridhara S, Dinesh KM, Vageesh TS, Nataraju SP. Heat use efficiency and helio-thermal units for maize genotypes as influenced by dates of sowing under southern transitional zone of Karnataka State. International Journal of Soil and Nature. 2011;2(3):529 -533.
- Cirilo AG, Andrade FH. Sowing date and maize productivity: I. Crop growth and dry matter partitioning. Crop Science. 1994;34 (4):1039-1043.
- Maddonni GA, Otegui ME, Bonhomme R. Grain yield components in maize: II. Postsilking growth and kernel weight. Field Crops Research. 2004;56(3):257-264.
- 26. Varma VS.. Standradisation of alternate planting window in hybrid seed production of maize (*Zea mays* L.) M. Sc. (Ag.) Thesis

Acharya NG. Ranga Agricultural University, Rajendranagar, Hyderabad; 2013.

- Shargi Y, Rad AHS, Band AA, Noormohammadi NG, Zahedi H. Yield and yield components of six canola (*Brassica napus* L.) cultivars affected by planting date and water deficit stress. African Journal of Biotechnology. 2011;10(46): 9309-9313.
- Dar NA, Singh KN, Ahmad L, Sofi JA, Bhat ME, Kotru R. Influence of dates of sowing, cultivars and different fertility levels on fodder oat (*Avena sativa* L.) under temperate conditions of Kashmir valley (India). Range Management and Agroforestry. 2014;35(1):51-55.
- 29. Darby HM, Lauer JG. Planting date and hybrid influence on corn forage yield and quality. Agronomy Journal .2002;94(2):281-289.
- Mokhtarpur H, Mosavat SA, Feyzbakhsh MT, Saberi A. Effect of sowing date and planting density on ear yield of sweet corn in summer sowing. Electronic Journal of Crop Production. 2008;1(1):101-113.
- 31. Salama ASH. Yield and nutritive value of maize (*Zea mays* L.) forage as affected by plant density, sowing date and age at harvest. Italian Journal of Agronomy. 2019;14(1383):114-121.
- 32. Gaile Z. Harvest time effect on yield and quality of maize (*Zea mays* L.) grown for silage. Latvian Journal of Agronomy. 2008;10:104-111.
- Ball DM, Collins M, Lacefield GD, Martin NP, Mertens DA, Olson KE, Putnam DH, Undersander, DJ, Wolf MW. Understanding forage quality. American Farm Bureau Federation Publication. 2001; 1(1):1-15.
- Koca OY, Canavar O. The effect of sowing date on yield and yield components and seed quality of corn (*Zea mays* L.). Scientific Papers Series A Agronomy. 2014;12(7):227-231.
- 35. Manjanagouda S, Sannagoudar, Lalitha BS, Prajwal Kumar GK, Prashant, Bhavya V. Effect of varieties, cutting, and nitrogen management on green fodder yield, nutrient uptake, available soil nutrient status, and economics of dual-purpose Pearl Millet (*Pennisetum glaucum* L.). International Journal of Current Microbiology and Applied Sciences. 2017;6 (12):214-226.
- 36. Somashekar KS, Shekara BG, Vidyashree DN, Lalitha BS, Bhavya V. Effect of

Different Seed Rates and Nitrogen Levels on Growth, Yield, Quality and Economics of Multi cut Fodder Sorghum (*Sorghum sudanense* L.). International Journal of Pure & Applied Bioscience. 2018; 6(1):1108-1115.

- Mahdi SS, Hasan B, Singh L. Influence of seed rate, nitrogen and zinc on fodder maize (*Zea mays* L.) in temperate conditions of western Himalayas. Indian Journal of Agronomy 2012;57(1):85-88.
- Iqbal J, Khan R, Wahid A, Sardar K, Khan N, Ali M, Hussain M, Ali W, Ali M, Ahmad R. Effect of nitrogen and zinc on maize (*Zea mays* L.) yield components and plant concentration. Advances in Environmental Biology. 2016;10(10):203-209.
- Ali S, Sahiba Malik MA, Hassan FU, Ansar M. Growth of rainfed fodder maize under different levels of nitrogen and phosphorus. Pakistan Journal of Agriculture Research. 2012;25(3):35-37.
- 40. Panwar D, Chouhan D, Singh D, Singh RP, Nepalia V. Performance of Fodder Maize (*Zea mays* L.) Under Varying Plant Densities and Fertility Levels. International Journal of Current Microbiology and Applied Sciences. 2020;11:255-260.
- 41. Eltelib HA, Hamad MA, Ali EE. The effect of nitrogen and phosphorus fertilization on growth, yield and quality of forage maize (*Zea mays* L.) Journal of Agronomy. 2006;5(3):515-18.
- 42. Budakli EC, Celik N, Bayram G. Yield and quality of forge maize as influenced by plant density and nitrogen rate. Turkish Journal of Field Crops. 2010;15(2):128-132.
- Aslam M, Iqbal A, Zamir MI, Mubeen M, Amin M. Effect of different nitrogen levels and seed rates on yield and quality of maize fodder. Crop and Environment. 2011;2(2): 47-51.
- 44. Patel SB, Patel AG, Chavda JN, Patel MA. Effect of time of sowing and nitrogen levels

on yield attributes, yield, and quality parameters of rabi forage maize (*Zea mays* L.). International Journal of Chemical Studies. 2017;5(5):868-871.

- 45. Meena S, Kumar S, Kumar R, Tokas J, Devi NU, Satpal. Response of springplanted fodder maize to nitrogen and phosphorus levels. Forage Research. 2021;46(4):363-368.
- 46. Kakol NB, Alagubdagi SC, Hosamani SV. Effect of Seed Rate and Nitrogen Levels on Forage yield and Quality of oat (*Avena sativa* L.). Indian Journal of Animal Nutrition. 2003;20(2):149-154.
- 47. Almodares A, Jafarinia M, Hadi MR. The effects of nitrogen fertilizer on chemical compositions in corn and sweet sorghum. American-Eurasian Journal of Agricultural and Environmental Sciences. 2009;6(4): 441-446.
- 48. Saruhan V, Sireli HD. An investegate on on the effect of plant densities and nitrogen doses on ear, stem and leaf yields of maize (*Zea Mays* L.). Journal of Agriculture 2005;9(2):45-53.
- 49. Chaudhary DP, Kumar A, Mandhania SS, Srivastava P, Kumar RS. Maize as fodder? An alternate approach. Technical bulletin, directorate of maize research, Indian Council of Agricultural Research, New Delhi; 2012.
- 50. Martin M, Williams II. Sweet corn growth and yield responses to planting dates of the North Central United States. Horticulture Science. 2008;43(6):1775-1779.
- Puri KP, Tiwana US. Effect of organic and inorganic sources of nitrogen in forage maize. Forage Research. 2008;34(1):62-63.
- 52. Saimaheswari K, Prathima T, Subramanyam D, Latha P. Agronomic evaluation of fodder sorghum varieties under different dates of sowing. Current Journal of Applied Science and Technology. 2020;39(9):25-32.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/119357