



Host Range and Distribution of Invasive Pest, Rugose Spiralling Whitefly *Aleurodicus rugioperculatus* Martin in Telangana, India

Islavath. Rahul Naik ^{a++*}, T. Pavani ^{b#}, S. Upendhar ^{ct}
and P. Neelima ^{d#}

^a Department of Entomology, PJTSAU, Hyderabad, India.

^b Department of Entomology, PJTSAU, Aswaraopeta, India.

^c Department of Entomology, PJTSAU, Rajendranagar, Hyderabad, India.

^d Department of Horticulture, PJTSAU, Aswaraopeta, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i121839>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129283>

Original Research Article

Received: 28/10/2024

Accepted: 30/12/2024

Published: 30/12/2024

ABSTRACT

Rugose Spiralling Whitefly (RSW), *Aleurodicus rugioperculatus*, is an invasive pest that invaded India in 2016, damaging coconut and other crops in Tamil Nadu, Kerala, Karnataka, and Andhra Pradesh. RSW stresses plants by depleting water and nutrients and promotes sooty mold growth

⁺⁺ Research Scholar;

[#] Assistant Professor;

[†] Associate Professor;

*Corresponding author: E-mail: islavath.rahul@gmail.com;

Cite as: Naik, Islavath. Rahul, T. Pavani, S. Upendhar, and P. Neelima. 2024. "Host Range and Distribution of Invasive Pest, Rugose Spiralling Whitefly *Aleurodicus rugioperculatus* Martin in Telangana, India". *Journal of Advances in Biology & Biotechnology* 27 (12):923-34. <https://doi.org/10.9734/jabb/2024/v27i121839>.

through honeydew excretion, reducing photosynthesis. The present study conducted, field surveys were conducted across major oil palm-growing districts of Telangana, including Bhadradi Kothagudem and Khammam, to assess the pest's distribution, damage intensity, infestation levels, and host range. This research provides critical insights for developing effective pest management strategies to mitigate the impact of RSW on economically significant crops.

Fifty host plants from 29 families were identified as hosts for RSW. Of these, 10 plants, including coconut, oil palm, banana, sapota, mango, and guava, supported all life stages of RSW, while the remaining host plants exhibited only the egg stage. The damage caused by RSW was particularly severe in coconut, oil palm, sapota, mango, and guava, highlighting their susceptibility. This comprehensive study underscores the varying levels of host suitability and damage intensity across different plant species, providing valuable insights into the pest's biology and its impact on key crops.

Keywords: *Plantation crops; fodder crops; vegetable crops; spices; avenue trees; field crops; ornamental fruits; medicinal crops; fruit crops; oil seed crops; Invasive pest.*

1. INTRODUCTION

The rugose spiralling whitefly (RSW), *Aleurodicus rugioperculatus* Martin, an invasive species, was first reported feeding on gumbo limbo (*Bursera simaruba*) in South Florida in 2004. In India, it was initially observed on coconut (*Cocos nucifera*) in Pollachi, Tamil Nadu, during August 2016 (Selvaraj et al., 2016; Sundharaj and Selvaraj, 2017). Subsequently, RSW was found feeding on banana, Indian almond, sapota, maize, oil palm, mango, cashew, custard apple, and various ornamental plants across Karnataka, Kerala, Andhra Pradesh, Telangana, Goa, Assam, and more recently in West Bengal, Lakshadweep, Maharashtra, Gujarat, Odisha, Chhattisgarh, and Meghalaya (Martin, 2004). Continuous feeding by nymphs and adults causes water and nutrient loss in host plants, while the release of honeydew on the upper surfaces of lower leaves promotes the growth of sooty mold. This sooty mold significantly reduces the plant's photosynthetic efficiency, further impacting its health and productivity (Capinera, 2008). In the Indian subcontinent, this invasive pest has spread to numerous locations, colonizing a wide range of host plants. Where previous studies reported infestation levels of 40–60% on coconut and 25–40% on banana. In several areas of Tamil Nadu and Kerala, the infestation was severe enough to cause complete drying of banana leaves (Selvaraj et al., 2016).

In India, six species known to infest coconut palms are *Aleurocanthus arecae* David and Manjunatha, *Aleurodicus dispersus* Russell, *Aleurodicus rugioperculatus* Martin, *Aleurotrachelus atratus* Hempel, *Paraleyrodes*

minei Iaccarino, and *Paraleyrodes bondari* Peracchi (Selvaraj et al., 2019).

Several natural enemies, such as *Encarsia guadeloupae* Viggiani (Hymenoptera: Aphelinidae), *Encarsia dispersa*, chrysopids, and coccinellids, were observed feeding on RSW (Elango and Nelson, 2020). A notable degree of shared parasitism was identified between coconut and intercrop herbivores, indicating that pest populations could be suppressed through indirect interactions. Given the rapid spread of RSW, it is essential to study the extent of damage across various agroclimatic zones in Telangana, identify its favourable hosts and natural enemies, and analyze its coexistence patterns. With these objectives, the present study was conducted to examine the distribution, host range, and status of the invasive rugose spiralling whitefly (*Aleurodicus rugioperculatus*) in Telangana.

2. MATERIALS AND METHODS

Identification of host range and estimation of extent of damage by *A. rugioperculatus* in Telangana: Using the scale and infestation index formula developed by Srinivasan et al. (2016), the identified host plants were surveyed and assessed to evaluate the extent of Rugose Spiralling Whitefly infestation across various host plants.

The infestation index for RSW was computed in coconut and oil palm based on the formula given below to categorize the coconut and oil palm gardens as low, medium and high intensity infestation on RSW (Srinivasan et al., 2016).

Table 1. Damage rating scale for the infestation of *A. rugioeperculatus*

No. of egg spirals /leaflet	Grade	Category	Infestation Index
No spirals – Sooty mold encrustation noticed	0	Nil	0.0
Less than 10 spirals/leaflet – Presence of Sooty mold encrustation in 5 to 6 lower fronds	1	low	0.01 -1.00
10 to 20 spirals/leaflet -Presence of Sooty mold encrustation in 10 to 12 fronds	2	Medium	1.01 – 2.00
More than 20 spirals/leaflet - Presence of Sooty mold encrustation in more than 12 fronds	3	High	2.01 - 3.00

Infestation index = ((No. of palms under scale 0 × 0) + (No of palms under scale 1 × 1) + (No of palms under scale 2 × 2) + (No of palms under scale 3 × 3) / Total no. of palms observed)

Hyophorbe lagenicaulis, Jatropha curcas, Saraca asoca, Acalypha wilkesiana, Dyspsis lutescens, Dracaena reflexa, Epipremnum aureum, Canna indica, Sterlizia reginae, Ixora coccinea, Aegel marmelos, Grewia asiatica, Citrus ssp, Annona reticulata, Aloe vera, Ficus religiosa and Mentha spicata.

Statistical analysis: The data was analyzed using analysis of variance (ANOVA) using AGRES 3.01 and AGDATA software. Data in the form of percentages were transformed to arcsine values and those in numbers were transformed to square root values where ever necessary and were analyzed. The mean values of different treatments were compared using DMRT at 5 per cent level of significance.

Among the recorded hosts, coconut (*Cocos nucifera*), oil palm (*Elaeis guineensis*), sapota (*Achras zapota*), Guava (*Pisidium guajava*) and Banana (*Musa paradisiaca*) were found to be heavily infested with *A. rugioeperculatus*, likely due to the pest's host preference, which is primarily driven by its nutritional needs. The infestation of RSW was also observed on the reproductive parts of banana, guava, mango and sapota, where egg spirals and sooty mold deposition were evident, as shown in Plate 4 respectively.

3. RESULTS AND DISCUSSION

Understanding the host range of a pest is crucial for developing an effective Integrated Pest Management (IPM) strategy. Given that *A. rugioeperculatus* is a relatively new invasive pest in India, and specifically in Telangana, the present study aims to identify its host range. This information will help assess the pest's potential impact, particularly on horticultural crops in the region.

Interestingly, some plant species such as castor, maize, banyan tree and sugarcane appeared to deter the complete development of RSW, though adult whiteflies were still observed feeding and laying eggs on these plants. The hosts listed in Table 2 and depicted in Plate 1 supported all life stages of RSW (from egg to adult). Among the identified hosts, plants from families like Arecaceae and Strelitziaceae were particularly favored by the pest for infestation.

In the present study, comprehensive surveys were conducted in the Bhadradi Kothagudem and Khammam districts of Telangana, providing valuable insights into the host range of *A. rugioeperculatus* (RSW). During the study period, *A. rugioeperculatus* was recorded on 50 plant species across 29 families, each harboring different life stages of the pest. Of these, 17 families were represented by a single species, while five families Poaceae, Asparagaceae, Strelitziaceae, Anacardiaceae and Annonaceae each hosted two species. Six families—Moraceae, Euphorbiaceae, Fabaceae, Malvaceae, Anacardiaceae and Annonaceae each hosted three species. Notably, 22 new host plants for RSW were identified: *Coccinia grandis, Curcumal longa, Magnolia champaca, Pongamia pinnata, Cassia angustifolia,*

Host that supported all the life stages are ivy gourd, banyan tree, champak, karanj, cassia, acalypha, butterfly palm, travellers tree, song of india, devil's ivy, oil palm, coconut, phalsa, sweet orange, mango, banana, sapota, guava, custard apple, jack fruit, indian almond, aloe and neem.

Host that supported both egg and adult stages are Drumstick, sugarcane, turmeric, castor, jatropha, bird of paradise and ixora.

Host that supported only egg stages are ivy gourd, drum stick, maize, sugar cane, banyan

Table 2. Crops identified in which all life stages of *A. rugioeperculatus* noticed in Telangana during 2022-2024

Sl.no	Common Name	Scientific Name	Order	Family	Life Stage			No. of egg spirals/ leaflet/leaf	Grade	Category
					Egg	Nymph	Adult			
Vegetable crops										
1	Ivy gourd	<i>Coccinia grandis</i>	Cucurbitales	Cucurbitaceae	+	+	+	<10	1	Low
2	Drum stick	<i>Moringa oleifera</i>	Brassicales	Moringaceae	+	-	+	<10	1	Low
Field crops										
1	Maize	<i>Zea mays</i>	Poales	Poaceae	+	-	-	<10	1	Low
2	Sugar cane	<i>Saccharum officinarum</i>	Poales	Poaceae	+	-	+	<10	1	Low
Forest tree										
1	Banyan tree	<i>Ficus benghalensis</i>	Rosales	Moraceae	+	+	+	<10	1	Low
Spices										
1	Turmeric	<i>Curcuma longa</i>	Zingiberales	Zingiberaceae	+	-	+	<10	1	Low
Oil seed crop										
1	Castor	<i>Ricinus communis</i>	Malpighiales	Euphorbiaceae	+	-	+	<10	1	Low
Avenue trees										
1	Champak	<i>Magnolia champaca</i>	Magnoliales	Magnoliaceae	+	+	+	10-20	2	Medium
2	Karanj	<i>Pongamia pinnata</i>	Fabales	Fabaceae	+	+	+	10-20	2	Medium
3	Cassia	<i>Cassia angustifolia</i>	Fabales	Fabaceae	+	+	+	10-20	2	Medium

Table 3. Crops identified in which all life stages of *A. rugioperculatus* noticed in Telangana during 2022-2024

Sl.no	Common Name	Scientific Name	Order	Family	Life Stage			No. of egg spirals /leaflet/leaf	Grade	Category
					Egg	Nymph	Adult			
Oranamental crops										
1.	Royal bottle palm	<i>Hyophorbe lagenicaulis</i>	Arecales	Arecaceae	+	-	-	<10	1	Low
2.	Jatropha	<i>Jatropha curcas</i>	Malpighiales	Euphorbiaceae	+	-	+	<10	1	Low
3.	Ashoka	<i>Saraca asoca</i>	Fabales	Fabaceae	+	-	-	<10	1	Low
4.	Acalypha	<i>Acalypha wilkesiana</i>	Malpighiales	Euphorbiaceae	+	+	+	<10	1	Low
5.	Butterfly palm	<i>Dypsis lutescens</i>	Arecales	Arecaceae	+	+	+	<10	1	Low
6.	Hibiscus	<i>Hibiscus rosa-sinensis</i>	Malvales	Malvaceae	+	-	-	<10	1	Low
7.	Spider planet	<i>Chlorophytum comosum</i>	Asparagales	Asparagaceae	+	-	-	<10	1	Low
8.	Travellers tree	<i>Ravenala madagascariensis</i>	Zingiberales	Strelitziaceae	+	+	+	10-20	2	Medium
9.	Song of India	<i>Dracaena reflexa</i>	Asparagales	Asparagaceae	+	+	+	<10	1	Low
10.	Devil's ivy	<i>Epipremnum aureum</i>	Alismatales	Araceae	+	+	+	<10	1	Low

Table 4. Crops identified in which all life stages of *A. rugioeperculatus* noticed in Telangana during 2022-2024

Sl.no	Common Name	Scientific Name	Order	Family	Life Stage			No. of egg spirals/ leaflet/leaf	Grade	Category
					Egg	Nymph	Adult			
Ornamental crops										
1	Canna	<i>Canna indica</i>	Zingiberales	Cannaceae	+	-	-	<10	1	Low
2	Bird of paradise	<i>Strelitzia reginae</i>	Zingiberales	Strelitziaceae	+	-	+	10-20	2	Medium
3.	Silver oak tree	<i>Grevillea robusta</i>	Proteales	Proteaceae	+	-	-	<10	1	Low
4.	Ixora	<i>Ixora coccinea</i>	Gentianales	Rubiaceae	+	-	+	<10	1	Low
Plantation crops										
1.	Areca nut	<i>Areca catechu</i>	Arecales	Areceae	+	-	-	<10	1	Low
2.	Cashew nut	<i>Anacardium occidentale</i>	Sapindales	Anacardiaceae	+	-	-	<10	1	Low
3.	Cocoa	<i>Theobroma cacao</i>	Malvales	Malvaceae,	+	-	-	<10	1	Low
4.	Oil palm	<i>Elaeis guineensis</i>	Arecales	Areceae	+	+	+	>20	3	High
5.	Coconut	<i>Cocos nucifera</i>	Arecales	Areceae	+	+	+	>20	3	High
Fodder crops										
1.	Napier grass	<i>Pennisetum purpureum</i>	Poales	Poaceae	+	-	-	<10	1	Low

Table 5. Crops identified in which all life stages of *A. rugioeperculatus* noticed in Telangana during 2022-2024

S. No.	Common Name	Scientific Name	Order	Family	Life Stage			No. of egg spirals/ leaflet/leaf	Grade	Category
					Egg	Nymph	Adult			
Fruit crops										
1.	Bael	<i>Aegle marmelos</i>	Sapindales	Rutaceae	+	-	-	<10	1	Low
2.	Phalsa	<i>Grewia asiatica</i>	Malvales	Malvaceae	+	+	+	<10	1	Low
3.	Sweet orange	<i>Citrus sinensis</i>	Sapindales	Rutaceae	+	+	+	<10	1	Low
4.	Citrus	<i>Citrus spp</i>	Sapindales	Rutaceae	+	-	-	<10	1	Low
5.	Water Rose apple	<i>Syzygium aqueum</i>	Myrtales	Myrtaceae	+	+	-	<10	1	Low
6.	Ramphal	<i>Annona reticulata</i>	Magnoliales	Annonaceae	+	-	-	<10	1	Low
7.	Mango	<i>Mangifera indica</i>	Sapindales	Anacardiaceae	+	+	+	<10	1	Low
8.	Banana	<i>Musa spp.</i>	Zingiberales	Musaceae	+	+	+	>20	3	High
9.	Sapota	<i>Manilkara zapota</i>	Ericales	Sapotaceae	+	+	+	Spirals on entire leaf	3	High
10.	Guava	<i>Psidium guajava</i>	Myrtales	Myrtaceae	+	+	+	>20	3	High

Table 6. Crops identified in which all life stages of *A. rugioeperculatus* noticed in Telangana during 2022-2024

S. No.	Common Name	Scientific Name	Order	Family	Life Stage			No. of egg spirals/ leaflet/leaf	Grade	Category
					Egg	Nymph	Adult			
Fruit crops										
1.	Custard apple	<i>Annona squamosa</i>	Magnoliales	Annonaceae	+	+	+	<10	1	Low
2.	Jack fruit	<i>Artocarpus heterophyllus</i>	Rosales	Moraceae	+	+	+	<10	1	Low
3.	Jamun	<i>Syzygium cumini</i>	Myrtales	Myrtaceae	+	-	-	<10	1	Low
4.	Indian Almond	<i>Terminalia catappa</i>	Myrtales	Combretaceae	+	+	+	<10	1	Low
Medicinal crop										
1.	Aloe	<i>Aloe vera</i>	Asparagales	Asphodelaceae	+	+	+	<10	1	Low
2.	Neem	<i>Azadirachta indica</i>	Sapindales	Meliaceae	+	+	+	<10	1	Low
3.	Camphor trees	<i>Cinnamomum camphora</i>	Laurales	Lauraceae	+	-	-	<10	1	Low
4.	Indian gooseberry	<i>Phyllanthus emblica</i>	Malpighiales	Phyllanthaceae	+	-	+	<10	1	Low
5.	Peppel	<i>Ficus religiosa</i>	Rosales	Moraceae	+	-	+	<10	1	Low
6.	Mint	<i>Mentha</i>	Lamiales	Lamiaceae	+	-	+	<10	1	Low

tree, turmeric, castor, champak, karanj, cassia, royal bottle palm, jatropha, ashoka, acalypha, butterfly palm, hibiscus, spider plant, travellers tree, song of india, devil's ivy, canna, bird of paradise, silver oak tree, ixora, areca nut, cashew nut, cocoa, oil palm, coconut, napier grass, bael, phalsa, sweet orange, citrus, water rose apple, ramphal, mango, banana, sapota, guava, custard apple, jack fruit, jamun, indian almond, aloe, neem, camphor tree, indian gooseberry, peppel and mint.

Plants supporting all life stages provide habitat and resources for eggs, larvae, pupae, and adults of specific organisms. However, some plants, like ivy gourd, support only the egg stage. Despite being listed in both categories, this distinction arises from the insect's selective use of hosts. Ivy gourd allows eggs to be laid but lacks the necessary resources (e.g., nutrition or shelter) for subsequent stages like larval feeding or pupation. Similarly, other hosts listed under the "egg stage only" category, such as drumstick, maize, and banyan tree, may serve as temporary surfaces or environments for egg-laying without facilitating full life cycle development.

During the study, it was observed that intercropping might be a primary factor contributing to the widespread prevalence of *A. rugioperculatus* (RSW) on multiple host plants. Intercropped plants initially serve as substrates for egg-laying and pest development. While early infestations were unstable, over several generations, the pest became well-established on these new host plants. Eventually, these hosts acted as sources of inoculum, facilitating the spread of the pest to other plants (Pasha & Muthuraju, 2023).

Intercropping facilitates the spread of *A. rugioperculatus* (RSW) by providing diverse plants for egg-laying and initial pest development. Over time, pests adapt and establish on these hosts, turning them into reservoirs of infestation. This creates a cycle where intercropped plants act as inoculum sources, accelerating pest dispersal to nearby crops (Pasha & Muthuraju, 2023).

The infestation of RSW was also observed in nurseries, orchards, horticultural gardens, and plantations along highways. When the surface characteristics of a new host plant differed from those of previous hosts, the pest appeared to modify its feeding habits to adapt to the new

host. This adaptation may be attributed to epigenetic changes in the pest, enabling it to survive on the new plant species.

Whittaker (1970) suggested that allelochemicals present in various plant species can significantly influence the growth and development of feeding insects. This may explain the higher incidence of *A. rugioperculatus* on certain crop families, such as *Arecaceae*, *Musaceae*, and *Myrtaceae*.

Mannion (2010) documented a wide range of host plants for RSW, including woody ornamentals, palms, and fruits. In Florida, between 2009 and 2012, 22% of the RSW-infested hosts were palm species, 16% were gumbo limbo, 10% were *Calophyllum sp.*, 9% were avocado, 4% were dark olive, and 3% were various mango varieties (Francis et al., 2016). Shanas et al. (2016) reported a total of 17 plant species from 11 families as preferred hosts of RSW in Kerala. Selvaraj et al. (2016) observed 25–40% leaf infestation on banana, with total leaf drying in several areas of Tamil Nadu and Kerala.

Srinivasan et al. (2016) recorded several preferred hosts, including coconut, sapota, custard apple, citrus, nutmeg, hibiscus, banana, okra, and guava, as well as alternate hosts such as parthenium, cassava, areca nut, and pepper. However, not all of these hosts supported the complete life cycle of the pest in Tamil Nadu.

Selvaraj et al. (2019) documented the incidence of RSW in West Bengal on a range of host plants, including coconut, banana, mango, sapota, Spanish cherry, hibiscus, guava, citrus, jamun, ficus, custard apple, areca nut, jackfruit, water apple, akashmoni, areca palm, and betel vine.

The findings of the present study are consistent with those of previous researchers, including Mannion (2010), Francis et al. (2016), Shanas et al. (2016), Selvaraj et al. (2016), Srinivasan et al. (2016), and Selvaraj et al. (2019).

In the present study, nine ornamental plants were observed that did not support all stages of *A. rugioperculatus* (RSW). This finding aligns with the observations of Krishnarao et al. (2019), who noted that while certain ornamental plants did not facilitate the complete development of RSW, adult whiteflies were still attracted to these plants for feeding and egg-laying. In field crops

like maize and sugarcane, only eggs were recorded, with no other developmental stages of the pest. This is consistent with the findings of Bhavani et al. (2020), who reported that while distinctive egg spirals were found on leaves, no nymphs or pupae were observed.

The results are in agreement with the findings of Elango and Nelson (2020) and Alagar et al. (2020), who also reported that neem and Indian almond supported all the stages of RSW when compared to the present study.




Plate 1. Host plants of *A. rugioeperculatus* A) Ivy gourd B) Drumstick C) Maize D) Sugar cane E) Banyan tree F) Travellers tree G) Castor H) Champak I) Karanj




Plate 2. Host plants of *A. rugioeperculatus* A) Cassia B) Royal bottle palm C) Jatropha D) Ashoka E) Acalypha F) Butterfly palm G) Jasmine H) Sugarcane I) Napier grass J) Custard apple




Plate 3. Host plants of *A. rugioperculatus* A) Jack fruit B) Spider planet C) Travellers tree D) Song of India E) Devils ivy F) Canna G) Bird of paradise H) Silver oak tree I) Ixora J) Areca nut




Plate 4. Host plants of *A. rugioperculatus* A) Cashew nut B) Cocoa C) Oil palm D) Coconut E) Bael F) Phalsa G) Sweet orange H) Water rose apple I) Ram phal J) Mango fruit

The study also recorded 22 new host plants, indicating that the host range of RSW is expanding, with new reports emerging from different areas. The spread of the pest appears to be influenced by the transport of planting material containing various developmental stages of the pest, as well as human activity. Additionally, prevailing winds and favourable climatic conditions play a significant role in the pest's dispersal to new areas. Given that

whiteflies can survive extreme weather conditions and have a high dispersal rate, effective management of this pest will require the adoption of an integrated pest management (IPM) strategy.

Drawbacks of present findings:

- **Limited Geographic Coverage:** The study was confined to specific districts in

Telangana, such as Bhadradi Kothagudem and Khammam, which may not represent the full range of agroclimatic conditions and host plants across the state or neighboring regions.

- **Lack of Long-Term Data:** The research provides a snapshot of RSW infestation but does not account for seasonal or inter-annual variations, which could influence pest dynamics, host preferences, and infestation intensity.
- **Insufficient Exploration of Biological Controls:** While natural enemies were mentioned, the study lacks detailed analysis of their efficacy, interactions with RSW, or practical recommendations for integrating them into pest management strategies.

4. CONCLUSION

The Rugose Spiralling Whitefly (RSW), *Aleurodicus rugioperculatus* Martin, is an invasive pest that first appeared in India in 2016, severely affecting crops like coconut, oil palm, banana, sapota, mango, and guava. RSW depletes plant water and nutrients and promotes sooty mold growth through honeydew secretion, significantly reducing photosynthetic efficiency.

A field study conducted across Telangana, particularly in Bhadradi Kothagudem and Khammam districts, identified 50 host plants from 29 families. While 10 crops, including coconut and oil palm, supported all life stages of RSW, other plants predominantly exhibited only the egg stage. Infestation was particularly severe in coconut and oil palm, with intercropping identified as a key factor in RSW's spread.

Infestation levels were classified into low, medium, and high categories based on egg spiral counts, and a damage rating scale was established. Several ornamental plants and field crops like maize and sugarcane only supported the egg stage, suggesting partial host suitability. Additionally, 22 new host plants were identified, indicating an expanding host range.

Natural enemies, including *Encarsia guadeloupae* and other parasitoids, were observed, showing potential for biological control. However, RSW's adaptability to new hosts and rapid dispersal due to favourable conditions highlight the urgency of adopting Integrated Pest Management (IPM) strategies.

This study emphasizes understanding host suitability, infestation patterns, and natural enemies to mitigate RSW's impact on economically significant crops in India. Effective management will require a combination of cultural practices, biological control, and targeted use of insecticides.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Alagar, M., Sivakumar, V., Chinnaduari, S., Saravanan, P.A., Srinivasan, T and Praneetha, S. 2020. Bio intensive management of invasive rugose spiralling whitefly in coconut. *Biotica Research Today*. 2(8): 768-771.
- Bhavani, B., Kishore Varma, P and Bharatha Lakshmi, M. 2020. First report of rugose spiralling whitefly, *Aleurodicus rugioperculatus* Martin, an invasive pest on sugarcane in Andhra Pradesh, India. *Journal of Entomology and Zoology Studies*. 8(6): 1993-1999.
- Capinera, J.L. (2008) *Encyclopedia of entomology*. 2nd edn. Springer, Netherlands, p 4346.
- Elango, K., and Nelson, S.J. 2020. Effect of host plants on the behaviour of rugose spiralling whitefly (*Aleurodicus rugioperculatus*) and their natural enemies. *Research Journal of Agricultural Sciences*. 11(1): 120-123.
- Francis, A.W., Stocks, I.C., Smith, T.R., Boughton, A.J., Mannion, C.M and Osborne, L.S. 2016. Host plants and natural enemies of Rugose spiralling whitefly (Hemiptera: Aleyrodidae) in Florida. *Florida Entomologist*. 99(1): 150-153.
- Krishnarao, G and ChalapathiRao, N.B.V. 2019. Surveillance and Eco-friendly management of new invasive alien pest, rugose spiralling whitefly, *Aleurodicus rugioperculatus* Martin: Inherent menace.

- Journal of Applied Zoological Research*. 30 (2): 148-158.
- Mannion, C. 2010. Rugose spiralling whitefly, a new whitefly in South Florida. *Tropical Research and Education Center, University of Florida*. 5(23):17-18.
- Martin, J. H. (2004). Whiteflies of Belize (Hemiptera: Aleyrodidae). Part 1 introduction and account of the subfamily Aleurodicinae Quaintance & baker. *Zootaxa*, 681(1), 1-119.
- Pasha, B. A., & Muthuraju, R. (2023). Characterization of Gut Bacterial Diversity and its Functional Role of Coconut Rugose Spiraling Whitefly *Aleurodicus rugioperculatus* Martin. *Mysore Journal of Agricultural Sciences*, 57(4).
- Selvaraj, K., Sundararaj, R and Sumalatha, B.V. 2019. Invasion of the palm infesting whitefly, *Aleurotrachelus atratus* Hempel (Hemiptera: Aleyrodidae) in the Oriental region. *Phytoparasitica*. 47(7): 327-332.
- Selvaraj, K., Sundararaj, R., Venkatesan, T., Ballal, C.R., Jalali, S.K., Gupta, A. Mrudula, H.K. (2016) Potential natural enemies of the invasive rugose spiralling whitefly, *Aleurodicus rugioperculatus* Martin in India. *J Biol Control*. 30(4):236–239.
- Shanas, S., Job, J., Joseph, T and Krishnan, G.A. 2016. First report of the invasive rugose spiralling whitefly, *Aleurodicus rugioperculatus* Martin (Hemiptera: Aleyrodidae) from the old world. *Entomon*. 41(4): 365-368.
- Srinivasan, T., Saravanan, P.A., Josephraj Kumar, A., Rajamanickam, K., Sridharan, S., David, P.M.M., Natarajan, N and Shoba, N. 2016. Invasion of the rugose spiralling whitefly, *Aleurodicus rugioperculatus* Martin (Hemiptera: Aleyrodidae) in Pollachi tract of Tamil Nadu, India. *Madras Agricultural Journal*. 103(10-12): 349-353.
- Sundararaj, R and Selvaraj, K. (2017) Invasion of rugose spiralling whitefly, *Aleurodicus rugioperculatus* Martin (Hemiptera: Aleyrodidae): a potential threat to coconut in India. *Phytoparasitica*. 45(1):71–74.
- Whittaker, R.H. 1970. The biochemical ecology of higher plants. *Chemical ecology*. 3: 43-70.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129283>