



Effect of Chitosan Biopolymer on Microbial Decay in Tomato (*Solanum lycopersicum* L.) Variety 'Dev' under Ambient Temperature

**Mahendra Meena ^{a*}, Kinjal Mondal ^b, Vinod Saharan ^b,
K. K. Meena ^a, N. K. Gupta ^c, O. P. Garhwal ^a, L. N. Bairwa ^d,
Shalini Pilania ^e, Mital ^b and Deepak Meena ^f**

^a Department of Horticulture, Rajasthan Agricultural Research Institute, SKNAU, Jobner, Rajasthan, India.

^b Department of Molecular Biology and Biotechnology, Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan, India.

^c Department of Plant Physiology, SKNAU, Jobner, Rajasthan, India.

^d Department of Horticulture, SKN College of Agriculture, SKNAU, Jobner, Rajasthan, India.

^e Department of Horticulture, Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan, India.

^f Department of Genetics and Plant Breeding, Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i41706

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97199>

Original Research Article

Received: 01/01/2023

Accepted: 04/03/2023

Published: 06/03/2023

*Corresponding author: E-mail: mahendra966795@gmail.com;

ABSTRACT

The current study sought to investigate the influence of chitosan biopolymer against microbial decay in tomato. In the present study, bulk-chitosan at different concentrations (0.01, 0.04, 0.08, 0.12, 0.16, 0.20% w/v) along with control were evaluated on tomato variety 'Dev' by dipping fruits for 6 min and stored at ambient temperature ($27^{\circ}\text{C}\pm 2$). Bulk-chitosan at 0.16% was found most effective to prevent microbial decay and maintain sensory evaluation from day 1 to days 21 as compared to control. Based on the aforementioned results and comparative evaluation of different doses of bulk-chitosan, we concluded that chitosan is very effective at less concentration and thus exert minimum chemical load on the treated tomatoes. Therefore, it may claim that chitosan biopolymer have potential to protects tomato against microbial decay.

Keywords: Chitosan; biopolymer; microbial decay; tomato.

1. INTRODUCTION

"Tomato (*Solanum lycopersicum* L.) is the second most important vegetable crop after potato [1], and it is a model plant for researchers, especially for those studying the growth, yield and quality of fleshy fruits" [2]. "Tomato is one of the most important foods for humanity. The post-harvest maintenance of food is configured as one of the objectives to be achieved in the sustainable development proposed by FAO in 2030. Tomato contains higher amounts of lycopene, a type of carotenoid with antioxidant properties which is beneficial in reducing the incidence of chronic diseases like cancer and many other cardiovascular disorders" [3]. India is the 2nd largest (11.5%) producer of tomato in world. According to FAOSTAT [1], "tomato was grown world-wide in an area of 5.16 MH with the production of 189.13 MT and productivity of 36.6 MT per hectare. In India it was grown in 0.84 MH with the production of 21.18 MT and productivity of 25.06 MT per hectare". "Chitosan is a linear amino polysaccharide of glucosamine and N-acetylglucosamine units and obtained by alkaline deacetylation of chitin extracted from the exoskeleton of crustaceans such as shrimps and crabs as well from the cell walls of some fungi" [4]. "It is being widely used in different fields ranging from medicine, tissue and bone engineering to food sector, cosmetics, textiles, pharmaceutical, biotechnology, paper industry and in waste-water treatment" [5]. In agriculture, chitosan biopolymer is an ideal antimicrobial agent in preservation of fruits and vegetables. Chitosan is generally used to prevent microbial diseases and decay of fruits and vegetables through its pre- and post-harvest treatments [6]. Chitosan is considered as a strong antimicrobial agent due to its positive surface charge which alters the microbial membrane structure that eventually leads to leakage of cellular fluids and

death [7]. The NH_2 group present in it is responsible for its bioactivity against microbial community through inhibition of gene expression and protein synthesis [8]. Chitosan has been used to prevent post-harvest losses in fruits and vegetables. It is also effective in reducing post-harvest diseases caused by various microbes like *Botrytis cinerea* [9,10], *Penicillium expansum* [11,12], *Alternaria alternata* [13], *Colletotrichum gloeosporioides* [14] and *Rhizopus stolonifer* [6] by inhibiting spore germination, germ tube elongation and mycelial growth of fungal phytopathogens. In yet another study on tomato, it was found that application of chitosan at different concentrations in the range 0.01 to 1% significantly inhibited the growth of *Botrytis cinerea* and *Penicillium expansum* as compared with untreated tomato. Chitosan exhibits direct fungitoxic activity as well as elicits the biochemical defense responses in fruit [11]. On the basis of above facts, in present study, we proposed chitosan formulation, and delve its bioactivity by testing its effect on tomato variety 'Dev'. Efforts have been made to study the chitosan formulation against microbial decay and sensory evaluation under ambient temperature upto 21 days.

2. MATERIALS AND METHODS

Chitosan preparation: In present study, bulk-chitosan (BCH) formulations were prepared by dissolving chitosan (Mol. Wt. 50,000–190,000 and 80% N-deacetylation; Sigma-Aldrich, St. Louis, MO, USA) into 1% glacial acetic acid to get final concentrations of 0.01, 0.04, 0.08, 0.12, 0.16 and 0.20% (w/v) with adjusting pH 5.1 with 1N NaOH. The chitosan formulations thus prepared were used to treat tomato fruits.

Experimental details: The experiment was conducted during (September to November)

2018-2019 at the Department of Horticulture and Department of Molecular Biology and Biotechnology, Rajasthan College of Agriculture, Maharana Pratap University of Agriculture and Technology, Udaipur. Tomato variety *Dev* grown at the University farm, Rajasthan College of Agriculture, MPUAT was used in present investigation. Tomatoes with similar size, firmness and colour were selected for the experiment. The fruits with visible decay and mechanical damage were abandoned for the study. Selected tomato fruits were dipped into different concentrations of chitosan formulation (0.01, 0.04, 0.08, 0.12, 0.16 and 0.20%, w/v in water) along with control (distilled water) for 6 min and stored at ambient temperature.

Fungal decay index: Mechanical cut (0.5 cm x 1.0 cm) was positioned on tomatoes for stimulation of microbial infection. "Microbial decay was visually examined, considering the level of fungal growth on the tomato surface in the scale ranging from 1 to 5, where 1 = normal (no decay on fruit surface), 2 = trace (up to 5 % of fruit surface decayed), 3 = slight (5–20 % of fruit surface decayed), 4 = moderate (20–50 % of fruit surface decayed) and 5 = severe (>50 % of fruit surface decayed)" (Babalar et al. 2007). Results were expressed as fungal decay index.

2.1 Overall Organoleptic Score

Overall acceptability of the samples was evaluated through the standard sensory evaluation techniques. The sensory attributes

such as taste, flavor and acceptability was rated by using five-point hedonic scale (9-Excellent, 7-Very good, 5-Good, 3-Fair, 1-Poor) as well as 11 members panel of judges (Post harvest experts).

2.2 Statistical Analysis

Statistical analysis was performed with JMP software version 12 [15] using Turkey Kramer HSD test. Each experiment was repeated twice wherein each treatment consisted of minimum three replicates having five tomatoes each.

3. RESULTS AND DISCUSSION

In the present study, solutions of bulk-chitosan at different concentrations (0.01, 0.04, 0.08, 0.12, 0.16 and 0.20%, w/v) were prepared in 1% acetic acid. The pH of solutions was adjusted to 5.5 to eliminate the acidic damage to tomato fruits. Microbial decay was visually inspected up to 21 days considering the extent of microbial infection on fruit surface. Decay rate increased with storage time due to microbial infection. The lowest decay (5% at scale of 2) was found in fruits treated with 0.08 and 0.16% bulk-chitosan. Microbial decay contributes up to ~70% losses in tomato and is, therefore, very crucial to control it during storage [16]. Bulk-chitosan at 2-4% concentrations considerably controlled gray mould in wound inoculated tomato fruits [10] whereas, at 0.5-1% concentrations significantly inhibited the growth of gray and blue moulds in tomato fruits stored for 21 days in cold storage [11]. Various concentrations of bulk chitosan




Fig. 1. Effect of chitosan formulation on microbial decay at room temperature of tomato. Each value is mean of triplicate. Error bars represents \pm SE




Fig. 2. Effect of of chitosan formulation on sensory evaluation at room temperature of tomato
Each value is mean of triplicate. Error bars represents \pm SE

considerably controlled decay of strawberry, pomegranate and table grapes during storage [17-20]. In our results, bulk chitosan at 0.08 and 0.16% concentrations significantly controlled tomato decay up to 21 days of storage (Fig. 1). Results obtained in present investigation are better as compared with previous findings as very low concentration of bulk-chitosan comprehensively controlled the decay up to 21 days of storage at ambient temperature [21]. Positively charged chitosan effectively degrades microbial cell wall and also boosts plant's immunity by enhancing defense enzymes activities [22,23]. Sensory evaluation is another important parameter responsible for acceptability of tomatoes by consumers. Gao et al. [20] studied that flavour of table grape significantly decreased after 15 days of storage in untreated fruits while with chitosan treatment, the sensory parameters were maintained up to the end of storage period. Our results showed that 0.16% bulk-chitosan was fairly effective to preserve color, texture, flavour and overall acceptability of tomato fruit (Fig. 2).

4. CONCLUSION

The available review of literature and results of the present investigation concluded that amongst the various treatments, 0.16% bulk-chitosan significant results in microbial decay and sensory evaluation as compared with control up to 21 days of storage at ambient temperature. Based on the aforementioned results and comparative

evaluation of different doses of bulk-chitosan, we concluded that chitosan are very effective at less concentration and thus exert minimum chemical load on the treated tomatoes. Therefore, it is possible to assert that chitosan biopolymer has the capacity to shield tomatoes against microbial degradation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Food and Agriculture Organization (FAO). FAOSTAT Statistical Database of the United Nations Food and Agriculture Organization (FAO) statistical division. Rome; 2021.
2. Bertin N, Génard M. Tomato quality as influenced by preharvest factors. *Sci Hortic*. 2018;233:264-76.
3. Viskelis P, Radzevicius A, Urbonaviciene D, Viskelis J, Karkleliene R, Bobinas C. Biochemical parameters in tomato fruits from different cultivars as functional foods for agricultural, industrial, and pharmaceutical uses. In: *Tech Open*, 10 ElShemy H, editor. *Plants for the future*. 2015; 60873:5772.
4. Sucharitha KV, Beulah AM, Ravikiran K. Effect of chitosan coating on storage stability of tomatoes (*Lycopersicon*

- esculentum Mill.). Int Food Res J. 2018; 25(1):93-9.
5. Jiang YM, Li YB. Effect of chitosan coating on postharvest life and quality of longan fruit. Food Chem. 2001;73(2):139-43.
6. Bautista-Baños S, Hernández-Lauzardo AN, Velázquez-del Valle MG, Hernández-López M, Ait Barka E, Bosquez-Molina E et al. Chitosan as a potential natural compound to control pre and postharvest diseases of horticultural commodities. Crop Prot. 2006;25(2):108-18.
7. Goy RC, Britto D, Assis OBG. A review of the antimicrobial activity of chitosan. Assis Polimeros. Cien Tecnol. 2009;19(3):241-7.
8. Kong M, Chen XG, Xing K, Park HJ. Antimicrobial properties of chitosan and mode of action: A state of the art review. Int J Food Microbiol. 2010;144(1):51-63.
9. Elmer PAG, Reglinski T. Biosuppression of *Botrytis cinerea* in grapes. Plant Pathol. 2006;55(2):155-77.
10. Badawy MEI, Rabea EI. Potential of the biopolymer chitosan with different molecular weights to control postharvest graymold of tomato fruit. Postharvest Biol Technol. 2009;51(1):110-7.
11. Liu J, Tian S, Meng X, Xu Y. Effects of chitosan on control of postharvest diseases and physiological responses of tomato fruit. Postharvest Biol Technol. 2007;44(3):300-6.
12. Yu T, Li HY, Zheng XD. Synergistic effect of chitosan and *Cryptococcus laurentii* on inhibition of *Penicillium expansum* infections. Int J Food Microbiol. 2007;114(3):261-6.
13. Domínguez I, Ferreres F, Pascual del Riquelme F, Font R, Gil MI. Influence of preharvest application of fungicides on the postharvest quality of tomato (*Solanum lycopersicum* L.). Postharvest Biol Technol. 2012;72:1-10.
14. Ramos-Garcia M, Bosquez-Molina E, Hernandez-Romano J, Zavala-Padilla G, Terres Rojas E, Alia-Tejacal I et al. Hernandez-L opez M, Bautista-Banos S. Use of chitosan-based edible coatings in combination with other natural compound, to control *Rhizopus stolonifer* and *Escherichia coli* DH5a in fresh tomatoes. Crop Protection. 2012;38:1-6.
15. SAS. JMP: User's guide. Version 14.2. Cary, NC: SAS Institute, Inc.; 2019.
16. Meena M, Pilia S, Pal A, Mandhan S, Bhushan B, Saharan V. Cu-chitosan nano-net improves keeping quality of tomato by modulating physio-biochemical responses. Sci Rep. 2020;10(1):1-11.
17. Hajji S, Younes I, Affes S, Boufi S, Nasri M. Optimization of the formulation of chitosan edible coating supplemented with carotenoproteins and their use for extending strawberries postharvest life. Food Hydrocoll. 2018;83:375-92.
18. Candir E, Ozdemir AE, Aksoy MC. Effects of chitosan coating and modified atmosphere packaging on postharvest quality and bioactive compounds of pomegranate fruit cv. 'Hicaznar'. Sci Hortic. 2018;235:235-43.
19. Feliziani E, Landi L, Romanazzi G. Preharvest treatments with chitosan and other alternatives to conventional fungicides to control postharvest decay of strawberry. Carbohydr Polym. 2015;132: 111-7.
20. Gao P, Zhu Z, Zhang P. Effects of chitosan-glucose complex coating on postharvest quality and shelf life of table grapes. Carbohydr Polym. 2013;95(1):371-8.
21. El Ghaouth AE, Ponnampalam R, Castaigne F, Arul J. Chitosan coating to extend the storage life of tomatoes. Hortscience. 1992;27(9):1016-8.
22. Bai RK, Huang MY, Jiang YY. Selective permeabilities of chitosan-acetic complex membrane and chitosan-polymer complex for oxygen and carbon dioxide. Polym Bull. 1988;20(1):83-8.
23. Butler BL, Vergano PJ, Testin RF, Bunn JM, Wiles JL. Mechanical and barrier properties of edible chitosan films as affected by composition and storage. J Food Sci. 1996;61(5):953-6.

© 2023 Meena et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97199>